

Deworming Protocol



Fecal egg counts (FEC) provide us with valuable information that help us evaluate the efficacy of de-wormers being used and determine the best deworming treatment plan for your horse. Spring and Fall are peak transmis-sion time, by doing a FEC during these times and following a deworming protocol we can minimize pasture contamination and prevent parasitic disease and resistance.

Adult LOW Shedder Treatment Plan

Below 500 EPG on Fecal Egg Test

	SPRING	FALL
	RUN A FECAL TEST	RUN A FECAL TEST
	DEWORMER TO USE:	DEWORMER TO USE:
•	Moxidectin (QUEST)	Ivermectin with Praziquan- tel
•	Ivermectin for minia-	
	ture horses	Zimecterin Gold





Foal Treatment Plan

Adult HIGH Shedder Treatment Plan

Above 500 EPG on Fecal Egg Test

MONTH	FECAL TEST?	DEWORMER TO USE
January	NO	Ivermectin
March	YES	Moxidectin (QUEST)Ivermectin for miniature horses
May	NO	Double dose of Pyrantel (STRONGID)
July	NO	Ivermectin with Praziquantel Zimecterin Gold
October	YES	Moxidectin (QUEST)Ivermectin for miniature horses

Age	Treatment Plan	Brand Name of Dewormer
2 Months	Fenbendazole (10mg/kg)	Panacur
3 Months	Pyrantel	Strongid
4 Months	Ivermectin	Zimecterin, Bimectin, Ivermectin Paste
4 ½ Months		Fecal Test!
6 Months	Pyrantel	Strongid
8 Months	Ivermectin with Pra- ziquantel	Zimecterin Gold
10 Months	Pyrantel	Strongid
12 Months	Moxidectin	Quest for 700 lbs and up & Panacur Power Pak for under 700 lbs